

TIT/HEP-267
 NUP-A-94-18
 hep-th/9410090
 October, 1994

QUANTUM DIFFUSION PROCESS WITH A COMPLEX WEIGHT ON THREE DIMENSIONAL LATTICE*

MASAKO ASANO

*Department of Physics, Tokyo Institute of Technology,
 Oh-okayama, Meguro, Tokyo 152, Japan*

CHIGAK ITOI

*Department of Physics,
 College of Science and Technology, Nihon University,
 Kanda Surugadai, Chiyoda, Tokyo 101, Japan*

and

SHIN-ICHI KOJIMA

*Department of Physics, Tokyo Institute of Technology,
 Oh-okayama, Meguro, Tokyo 152, Japan*

Abstract

A rigorous definition of a path integral for a spinning particle in three dimensions is given on a regular cubic lattice. The critical diffusion constant and the associated critical exponents in each spin are calculated. Continuum field theories such as Klein-Gordon, Dirac and massive Chern-Simons theories are constructed near these critical points. The universality of obtained results is argued on some other lattices.

1 Introduction

A path integral representation of spinning particles in three dimensional Euclidian space is studied extensively [1] - [5]. The free energy of spinning particles with spin $J \in \mathbf{Z}/2$ can be expressed in terms of a bosonic path integral. It is well-known that the critical behavior of this model differs from that in Brownian motion. Since a path has to pay energy to bend itself due to the spin factor, a typical path in this model is much smoother than that in Brownian motion. And then, the spin factor affects the critical exponents of the model. This path integral representation has been used to argue a spinor field coupled to Chern-Simons field in [1] [4] [5]. However, it is difficult to check the self-consistency among several heuristic limitation procedures employed to

* An updated version of the talk presented by C. Itoi in The International Seminar Devoted to the 140th Birthday of Henri Poincaré IHPE Protovino, Russia, 27/June - 1/July, 1994 to appear in the Proceedings

define the path integral, and therefore the reliability of the nonperturbative consequence obtained by this method remains unclear.

In this talk, we report our recent work on construction of continuum spinor field theories in terms of random walks on a three dimensional lattice [6] [7]. The random walk is defined on lattice and the continuum limit is discussed rigorously. This model corresponds to a simple generalization of the Markovian process for solving two dimensional Ising model [9] to that in three dimensions with other spins. In the case of two dimensional Ising model, the number of diagrams in the high temperature expansion is evaluated in terms of a particular Markovian process. We find a critical point for each value of J and study its critical behaviors to construct continuum field theories. In regular cubic lattice, Klein-Gordon, Dirac and massive Chern-Simons fields can be constructed at corresponding critical points. In other lattice such as triclinic lattice, only Klein-Gordon and Dirac fields can be constructed. Therefore, the construction of the Klein-Gordon and the Dirac theories by this random walk model is expected to be universal but that of the massive Chern-Simons theory is not universal.

2 Definition of the model

Let us consider the following Markovian diffusion process on a three dimensional lattice with an amplitude depending on the preceding step. To define the free energy, we evaluate a sum over all paths leaving and returning to the origin. First, we work on regular cubic lattice. Let $\mathbf{x}$ and $\mathbf{e}_\alpha$ be a lattice site and a bond indices, respectively. The length of each bond is taken to be unity. We define a probability amplitude $A_t^{\alpha\beta}(\mathbf{x})$ for a particle that starts at the origin, moving in direction $\mathbf{e}_\alpha$ to propagate in t steps to a site $\mathbf{x}$ moving in direction $\mathbf{e}_\beta$. Here, $A_t^{\alpha\beta}(\mathbf{x})$ satisfies a recursion relation

$$A_{t+1}^{\alpha\beta}(\mathbf{x}) = \sum_{\gamma \neq -\beta} \kappa e^{iJS(\mathbf{e}_3, \mathbf{e}_\beta, \mathbf{e}_\gamma)} A_t^{\alpha\gamma}(\mathbf{x} - \mathbf{e}_\beta), \quad (1)$$

where κ is a diffusion constant and J is a spin parameter. The sum is taken over all nearest neighbor bonds $\mathbf{e}_\gamma$ at the site $\mathbf{x}$ except $\mathbf{e}_{-\beta}$, where we denote $\mathbf{e}_{-\gamma} = -\mathbf{e}_\gamma$. In our model any backtracking process is forbidden. $S(\mathbf{e}_\alpha, \mathbf{e}_\beta, \mathbf{e}_\gamma)$ is the area of a spherical triangle with vertices $\mathbf{e}_\alpha$, $\mathbf{e}_\beta$ and $\mathbf{e}_\gamma$ having an orientation, basically. Some ambiguous cases are defined by $S(\mathbf{e}_3, \mathbf{e}_{-3}, \mathbf{e}_1) = 0$, $S(\mathbf{e}_3, \mathbf{e}_{-3}, \mathbf{e}_{-1}) = 2\pi$ and $S(\mathbf{e}_3, \mathbf{e}_{-3}, \mathbf{e}_{\pm 2}) = \pm\pi$. The initial condition is given by

$$A_0^{\alpha\beta}(\mathbf{x}) = \delta^{\alpha\beta} \delta_{x_1 0} \delta_{x_2 0} \delta_{x_3 0}. \quad (2)$$

Here, we consider only half integer J . This diffusion process in two dimensions with $J = \frac{1}{2}$ enables us to calculate all orders in the high temperature expansion for the Ising model. In this representation, Majorana spinor is obtained as a critical theory of the two dimensional Ising model. Our three dimensional diffusion model is regarded as its generalization. A free energy

$f(\kappa, J)$ per unit volume is

$$f(\kappa, J) = -(-1)^{2J} \sum_{t=1}^{\infty} \sum_{\alpha=\pm 1}^{\pm 3} \frac{1}{t} A_t^{\alpha\alpha}(\mathbf{0}). \quad (3)$$

In momentum space we obtain the Fourier transformed recursion relation as a block diagonal form

$$B_{t+1}^{\alpha\beta}(\mathbf{p}) = \sum_{\gamma} \kappa B_t^{\alpha\gamma}(\mathbf{p}) Q^{\gamma\beta}(\mathbf{p}, J), \quad (4)$$

where $B_t^{\alpha\beta}(\mathbf{p})$ is the Fourier transform of the amplitude $A_t^{\alpha\beta}(\mathbf{x})$. The matrix $Q(\mathbf{p}, J)$ has the form

$$Q(\mathbf{p}, J) = \begin{bmatrix} e^{-ip_2} & e^{ip_1+i\frac{\pi}{2}J} & 0 & e^{-ip_1-i\frac{\pi}{2}J} & e^{-ip_3} & e^{ip_3-i\pi J} \\ e^{-ip_2-i\frac{\pi}{2}J} & e^{ip_1} & e^{ip_2+i\frac{\pi}{2}J} & 0 & e^{-ip_3} & e^{ip_3-i2\pi J} \\ 0 & e^{ip_1-i\frac{\pi}{2}J} & e^{ip_2} & e^{-ip_1+i\frac{\pi}{2}J} & e^{-ip_3} & e^{ip_3+i\pi J} \\ e^{-ip_2+i\frac{\pi}{2}J} & 0 & e^{ip_2-i\frac{\pi}{2}J} & e^{-ip_1} & e^{-ip_3} & e^{ip_3} \\ e^{-ip_2} & e^{ip_1} & e^{ip_2} & e^{-ip_1} & e^{-ip_3} & 0 \\ e^{-ip_2+i\pi J} & e^{ip_1+i2\pi J} & e^{ip_2-i\pi J} & e^{-ip_1} & 0 & e^{ip_3} \end{bmatrix} \quad (5)$$

The solution of the recursion equation (4) with (2) is

$$B_t(\mathbf{p}) = \{\kappa Q(\mathbf{p}, J)\}^t. \quad (6)$$

The free energy can be written in terms of the matrix $Q(\mathbf{p}, J)$

$$f(\kappa, J) = -(-1)^{2J} \int \frac{d^3p}{(2\pi)^3} \text{Tr} \sum_{t=1}^{\infty} \frac{1}{t} \{\kappa Q(\mathbf{p}, J)\}^t \quad (7)$$

This summation converges only when κ^{-1} is larger than the maximal absolute value of the eigenvalue of the matrix $Q(\mathbf{p}, J)$. This can be checked numerically. In this case, the summation becomes

$$f(\kappa, J) = (-1)^{2J} \int \frac{d^3p}{(2\pi)^3} \text{Tr} \log[1 - \kappa Q(\mathbf{p}, J)] \quad (8)$$

3 Critical Theories

Physical quantities become non-analytic at the critical point as a function of the diffusion constant κ , due to the finite convergence radius of the summation (7). If the correlation length diverges at the critical point, we can take a continuum field theory limit. The model at several critical points has a certain continuum limit. One can check the maximal absolute value in the eigenvalues of the matrix $Q(\mathbf{p}, J)$ is given at $\mathbf{p} = \mathbf{0}$ at least numerically. A critical diffusion constant κ_c is determined by the eigenvalue equation of $Q(\mathbf{0}, J)$

$$\det[\kappa_c^{-1} - Q(\mathbf{0}, J)] = 0. \quad (9)$$

At $\kappa = \kappa_c$, the free energy becomes non-analytic. Since $S(\mathbf{e}_\alpha, \mathbf{e}_\beta, \mathbf{e}_\gamma)$ in (1) is quantized by $\frac{\pi}{2}$ in our regular cubic lattice, the eigenvalue equation possesses the symmetry $J \rightarrow 4 + J$ and $J \rightarrow -J$. This is because that J is a half-integer and $Q(\mathbf{0}, -J)$ is equal to the transposed matrix

<i>Spin Factor</i>	<i>Eigenvalues</i>					
J=0	5	1	1	1	-1	-1
J=1/2	$1 + 2\sqrt{2}$	$1 + 2\sqrt{2}$	$1 - \sqrt{2}$	$1 - \sqrt{2}$	$1 - \sqrt{2}$	$1 - \sqrt{2}$
J=1	3	3	3	-1	-1	-1
J=3/2	$1 + \sqrt{2}$	$1 + \sqrt{2}$	$1 + \sqrt{2}$	$1 + \sqrt{2}$	$1 - 2\sqrt{2}$	$1 - 2\sqrt{2}$
J=2	3	3	1	1	1	-3

Table 1: Table of eigenvalues of the matrix $Q(\mathbf{0}, J)$

of $Q(\mathbf{0}, J)$. Thus we only have to consider the cases : $J = 0, \frac{1}{2}, 1, \frac{3}{2}, 2$. For each J , we find solutions of (9) as in Table 3 .

One can see several degeneracies. At n -th degenerate critical point, the amplitude is expressed as a linear combination of n independent eigenvectors and can survive as an excitation with long range correlation. Only $(J, \kappa_c^{-1}) = (0, 5), (\frac{1}{2}, 1 + 2\sqrt{2}), (1, 3), (\frac{3}{2}, 1 + \sqrt{2}), (2, -3)$ and $(2, 3)$ have to be concerned for the convergence of the sum (7). One can construct rotationally invariant field theories at critical points $(J, \kappa_c^{-1}) = (0, 5), (\frac{1}{2}, 1 + 2\sqrt{2}), (1, 3)$ and $(2, -3)$, if one tunes the diffusion constant κ to these critical points κ_c . Those with single, double and triple degeneracies give us Klein-Gordon, Dirac and massive Chern-Simons fields, respectively. We do not find any other critical points that give continuum field theory.

First, we show the behavior of the free energy at critical points which give field theories. At $(J, \kappa_c^{-1}) = (0, 5)$ and $(2, -3)$, the critical behaviors belong to the same universality class. The free energy around $(0, 5)$ is given

$$f(\kappa, 0) \simeq \int \frac{d^3 p}{(2\pi)^3} \log[\mathbf{p}^2 + 4s + O(p^4)], \quad (10)$$

where we introduce small parameter s

$$s = \frac{\kappa_c - \kappa}{\kappa_c} > 0.$$

This critical behavior belongs to the same universality class of the Brownian motion. The correlation length diverges at the critical point as $\sim s^{-\frac{1}{2}}$, thus the critical exponent $\nu = \frac{1}{2}$. This critical point gives the Klein-Gordon theory as the continuum limit. The free energy near these two critical points behave as

$$f(\kappa) \sim s^{\frac{3}{2}} \log s.$$

This indicates that the order of the phase transition is second and the critical exponent $\alpha = \frac{1}{2}$.

At $(J, \kappa^{-1}) = (\frac{1}{2}, 1 + 2\sqrt{2})$, the free energy is

$$f(\kappa, \frac{1}{2}) \simeq - \int \frac{d^3 p}{(2\pi)^3} \log[\mathbf{p}^2 + s^2 + O(p^4)] \sim s^3 \log s. \quad (11)$$

The correlation length becomes s^{-1} , and then $\nu = 1$ and $\alpha = -1$. The two component Dirac field is obtained as the continuum limit at this critical point :

$$f(\kappa, \frac{1}{2}) \simeq - \int \frac{d^3 p}{(2\pi)^3} \text{Tr} \log[-i\mathbf{p} \cdot \sigma - s], \quad (12)$$

where σ is the Pauli matrix used as the gamma matrix in three dimensions. Other four components correspond to heavy particles which does not affect critical phenomena.

Here, we comment on a problem with fermion doubling on a lattice. For $(J, \kappa^{-1}) = (\frac{1}{2}, 1 + 2\sqrt{2})$, the matrix (5) has the only one massless excitation at $\mathbf{p} = \mathbf{0}$. The constant κ^{-1} should be tuned at $-1 - 2\sqrt{2}$ but not at $1 + 2\sqrt{2}$ to obtain massless excitation at $\mathbf{p} = (\pi, \pi, \pi)$. The corresponding doubler is too heavy at $\kappa^{-1} = 1 + 2\sqrt{2}$ to propagate, as in the case of the Wilson fermion. Therefore, no fermion doubling exists in our random walk model.

At $(J, \kappa^{-1}) = (1, 3)$, the non-analytic part of the free energy at $\kappa = 1 + \frac{8}{3}$ is evaluated as follows:

$$f(\kappa, -1) \simeq \int \frac{d^3 p}{(2\pi)^3} \log[\frac{16}{27}s\mathbf{p}^2 + \frac{64}{27}s^3 + O(p^4)] \sim s^3 \log s. \quad (13)$$

The free energy can be written also in terms of the following massive vector field

$$f(\kappa, -1) \simeq \int \frac{d^3 p}{(2\pi)^3} \text{Tr} \log[i\mathbf{p} \cdot \mathbf{L} + 4s], \quad (14)$$

where the matrix $\mathbf{L}$ is a spin matrix with the magnitude 1. One can take $(L_\mu)_{\nu\lambda} = i\epsilon_{\mu\nu\lambda}$. Two critical exponents $\alpha = -1$ and $\nu = 1$ are same as those of the Dirac field. The continuum limit is taken practically by $s \rightarrow 0$ after the replacement $\mathbf{p} \rightarrow s^\nu \mathbf{p}$. The model has infinite ultraviolet cutoff by this procedure.

Our method can construct the propagator $[1 - \kappa Q(J, \mathbf{p})]^{-1}$, as well as the free energy. Then, the Lagrangians of three types of rotationally invariant field theories constructed above are given for the Klein-Gordon field

$$\mathcal{L}_0 = \partial_\mu \phi^* \partial_\mu \phi + m \phi^* \phi, \quad (15)$$

the Dirac field

$$\mathcal{L}_{\frac{1}{2}} = \bar{\psi}(-i\gamma_\mu \partial_\mu + m)\psi, \quad (16)$$

and the massive U(1) Chern-Simons field

$$\mathcal{L}_1 = -i\epsilon^{\mu\nu\lambda} A_\mu^* \partial_\nu A_\lambda + m A_\mu^* A_\mu. \quad (17)$$

Above these three types of critical theories agree with the result given by the continuum path integral [1]-[5].

The order of phase transition at $(J, \kappa^{-1}) = (\frac{3}{2}, 1 + \sqrt{2})$, and $(2, 3)$ seems higher than the second and then the continuum limit might exist at those critical points. One might expect spin $\frac{3}{2}$ field theory at $(\frac{3}{2}, 1 + \sqrt{2})$. Nevertheless, these continuum theories at $J = \frac{3}{2}$ and 2 possess no rotational symmetry. For example, the free energy at $(J, \kappa^{-1}) = (\frac{3}{2}, (1 + \sqrt{2})(1 + s))$ becomes

$$f(\kappa, \frac{3}{2}) \simeq - \int \frac{d^3p}{(2\pi)^3} \log[18(3 - 2\sqrt{2})\{s^2 \mathbf{p}^2 + \frac{s^4}{2} + \frac{1}{4}(p_2^2 p_3^2 + p_3^2 p_1^2 + p_1^2 p_2^2) + O(p^6)\}]. \quad (18)$$

This shows no rotational invariance, even though one takes the limit $s \rightarrow 0$. Thus, the result for a spin higher than 1 does not agree with that obtained in the continuum approach. We can conclude that the continuum random walk method derive wrong results for higher spin fields.

4 Summary and Discussions

We studied a random walk model with a spin factor on a three dimensional lattice. Relativistic field theories of Klein-Gordon, Dirac and massive Chern-Simons fields have been constructed by taking continuum limit near critical points. It is found that there is no fermion doubling in this model. On the other hand, however, field theory with a spin higher than 1 have not been obtained as a rotationally invariant model.

To check the universality of our results, we have to work on some other lattice structures. Here, we summarize the results we have obtained so far in some other lattices such as triclinic and body centered cubic lattices [7]. All obtained results in a regular cubic lattice are derived in a body centered cubic lattice as well. In any lattice we have examined, the Klein-Gordon and the Dirac fields are obtained always. However, the massive Chern-Simons field cannot be obtained in triclinic lattices. The triple degeneracy of the matrix $Q(\mathbf{0}, 1)$ is split up in such lattice. This fact implies that the construction of spin 0 and spin 1/2 fields seems universal, while that of spin 1 field is not universal.

Acknowledgement

The authors thank T. Hara, N. Sakai and G. Semenoff for critical comments and helpful discussions.

They are also grateful to J. Ambjørn, P. Kurzepa and P. Orland for notifying them of references.

References

- [1] A.M. Polyakov, Modn. Phys. Lett. **A3**, 325 (1988);
- [2] J. Ambjørn, B. Durhuus and T. Jonsson, J. Phys. A **21** (1988) 981;
Nucl. Phys. **330**(1990)509.
- [3] A. Yu. Alekseev and S. L. Shatashvili, Modn. Phys. Lett. **A3**(1988)1551;
P. B. Weigmann, Nucl. Phys. **B323** (1989) 311;
T. Jacobson, Phys. Lett. **B216** (1989) 150;
P. Orland, Int. J. Modn. Phys. **A4** (1989) 3615;
T. Jaroszewicz and P. Kurzepa, Ann. Phys. **210** (1991) 255.
- [4] S. Iso, C. Itoi, and H. Mukaida, Phys.Lett.**B236**(1990)287;
Nucl. Phys. **B346**(1990)293;
N. Shaju, R. Shankar, and M. Sivakumar, Modn. Phys. Lett. **A5**, (1990)593;
W. Chen and C. Itoi, Phys. Rev. Letts, **72** (1994)2527; *Spin Transmutation in (2+1) Dimensions* IFT-488-UNC, NUP-A-94-3, hep-th/9404107.
- [5] C.-H. Tze, Int. J. Modn. Phys. **A3** (1988)1959;
- [6] C. Itoi *Smooth Paths on Three Dimensional Lattice* NUP-A-94-9, hep-th/9406123.
- [7] M. Asano, C. Itoi and S. Kojima *Random Walk Construction of Spinor Fields on Three Dimensional Lattice* TIT/HEP-271, NUP-A-94-19
- [8] D. Brydges, J. Fröhlich and T. Spencer Commun. Math. Phys. **83** (1982) 123 ;
R. Fernandez, J. Fröhlich and A. Sokal *Random Walks, Critical Phenomena, and Triviality in Quantum Field Theory* (Springer-Verlag, Berlin Heidelberg, 1992)
- [9] S. Sherman, J. Math. Phys. **1** (1960) 202; **4** (1963) 1213.
N. V. Vdovichenko, JETP **20**(1965)477;
M. L. Glasser, Am. J. Phys. **38** (1970) 1033.